


AMENDMENTS TO THE CLAIMS

1. (Currently amended) A compound ~~having~~ comprising the structural formula I, stereoisomers, tautomers, racemics, prodrugs, metabolites thereof, or a pharmaceutically acceptable salt and/or solvate thereof,



formula I

wherein X_1 , X_2 , R_1 and R_2 are independently selected from the group ~~comprising~~ consisting of oxo, hydrogen, hydroxyl, oxyalkyl, alkyl, alkenyl, alkynyl, alkyloxy, alkyloxyalkyl, alkylthioalkyl, alkoxycarbonyl, alkylthiocarbonyl, alkanoyl, cycloalkylalkyl, cycloalkylcarbonyl, cycloalkylalkanoyl, cycloalkylthiocarbonyl, cycloalkylalkoxycarbonyl, cycloalkylalkoxythiocarbonyl, cycloalkylthioalkyl, alkylcarbonyloxyalkyl, cycloalkylcarbonyloxyalkyl, silyloxyalkyl, aralkyl, arylalkenyl, arylcarbonyl, aryloxy, aryloxyalkyl, arylthiocarbonyl, aralkoxycarbonyl, arylalkylthiocarbonyl, aryloxyalkyl, arylthioalkyl, haloalkyl, hydroxyalkyl, aralkanoyl, aroyl, aryloxyalkyl, aryloxyalkanoyl, carboxyl, alkenylcarbonyl, alkynylcarbonyl, Het¹, Het¹alkyl, Het¹oxyalkyl, Het¹aryl, Het¹aralkyl, Het¹cycloalkyl, Het¹alkoxycarbonyl, Het¹alkylthiocarbonyl, Het¹oxycarbonyl, Het¹thiocarbonyl, Het¹alkanoyl, Het¹aralkanoyl, Het¹aryloxyalkyl, Het¹alkyloxyalkyl, Het¹arylthioalkyl, Het¹aryloxyalkyl, Het¹aralkoxycarbonyl, Het¹aroyl, Het¹oxyalkylcarbonyl, Het¹alkyloxyalkylcarbonyl, Het¹aryloxyalkylcarbonyl, Het¹carbonyloxyalkyl, Het¹alkylcarbonyloxyalkyl, Het¹aralkylcarbonyloxyalkyl, Het²alkyl, Het²oxyalkyl, Het²alkyloxyalkyl, Het²aralkyl, Het²carbonyl, Het²oxycarbonyl, Het²thiocarbonyl, Het²alkanoyl, Het²alkylthiocarbonyl, Het²alkoxycarbonyl, Het²aralkanoyl, Het²aralkoxycarbonyl, Het²aryloxyalkyl, Het²aroyl, Het²aryloxyalkyl, Het²arylthioalkyl, Het²oxyalkylcarbonyl, Het²alkyloxyalkylcarbonyl, Het²aryloxyalkylcarbonyl, Het²carbonyloxyalkyl,

Het²alkylcarbonyloxyalkyl, Het²aralkylcarbonyloxyalkyl, cyano, CR³=NR⁴, CR³=N(OR⁴), aminocarbonyl, aminoalkanoyl, aminoalkyl, optionally substituted by one or more substituents independently selected from the group ~~comprising~~consisting of alkyl, aralkyl, aryl, Het¹, Het², cycloalkyl, alkyloxy, alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(alkyl)aminocarbonyl, aminosulfonyl, alkylS(=O)_t, hydroxy, cyano, halogen or amino optionally mono- or disubstituted wherein the substituents are independently selected from the group ~~comprising~~consisting of alkyl, aryl, aralkyl, aryloxy, arylamino, arylthio, aryloxyalkyl, arylaminoalkyl, aralkoxy, alkylthio, alkoxy, aryloxyalkoxy, ~~arylaminealkoxy~~arylaminealkoxy, aralkylamino, aryloxyalkylamino, arylaminoalkylamino, arylthioalkoxy, arylthioalkylamino, aralkylthio, aryloxyalkylthio, arylaminoalkylthio, arylthioalkylthio, alkylamino, cycloalkyl, cycloalkylalkyl, Het¹, Het², Het¹alkyl, Het²alkyl, Het¹amino, Het²amino, Het¹alkylamino, Het²alkylamino, Het¹thio, Het²thio, Het¹alkylthio, Het²alkylthio, Het¹oxy and Het²oxy, OR³, SR³, SO₂NR³R⁴, SO₂N(OH)R³, CN, CR³=NR⁴, S(O)R³, SO₂R³, CR³=N(OR⁴), N₃, NO₂, NR³R⁴, N(OH)R³, C(O)R³, C(S)R³, CO₂R³, C(O)SR³, C(O)NR³R⁴, C(S)NR³R⁴, C(O)N(OH)R⁴, C(S)N(OH)R³, NR³C(O)R⁴, NR³C(S)R⁴, N(OH)C(O)R⁴, N(OH)C(S)R³, NR³CO₂R⁴, NR³C(O)NR⁴R⁵, and NR³C(S)NR⁴R⁵, N(OH)CO₂R³, NR³C(O)SR⁴, N(OH)C(O)NR³R⁴, N(OH)C(S)NR³R⁴, NR³C(O)N(OH)R⁴, NR³C(S)N(OH)R⁴, NR³SO₂R⁴, NHSO₂NR³R⁴, NR³SO₂NHR⁴, P(O)(OR³)(OR⁴), wherein t is an integer between 1 and 2 and R³, R⁴ and R⁵ are each independently selected from the group ~~comprising~~consisting of hydrogen, hydroxyl, alkyl, alkenyl, alkynyl, aminoalkyl, aminoaryl, alkylcarbonylamino, arylcarbonylamino, alkylthiocarbonylamino and arylthiocarbonylamino;

wherein X₃ participates together with X₃' ~~to~~in an oxo functional group, or wherein X₃ and X₃' are independently selected from the group ~~comprising~~consisting of hydrogen, hydroxyl, sulfur, oxyalkyl, oxycarbonyl, alkyl, Het¹alkyl, alkyloxycarbonyl, alkenyl, alkynyl, aminoalkyl, aminoacyl, alkylcarbonylamino, alkylthiocarbonylamino, Het¹, glycosyl, thio derivatives thereof, carboxy derivatives thereof, amino derivatives thereof, amido derivatives thereof, hydroxyl-protected derivatives thereof, optionally substituted by one or more substituents independently selected from the group ~~comprising~~consisting of alkyl, aralkyl, aryl, Het¹, Het², cycloalkyl, alkyloxy,

alkyloxycarbonyl, carboxyl, aminocarbonyl; mono- or di(alkyl)aminocarbonyl, aminosulfonyl, alkylS(=O)_t, hydroxy, cyano, halogen or amino optionally mono- or disubstituted wherein the substituents are independently selected from the group comprising consisting of alkyl, aryl, aralkyl, aryloxy, arylamino, arylthio, aryloxyalkyl, arylaminoalkyl, aralkoxy, alkylthio, alkoxy, aryloxyalkoxy, ~~arylaminealkoxy~~ arylaminoalkoxy, aralkylamino, aryloxyalkylamino, arylaminoalkylamino, arylthioalkoxy, arylthioalkylamino, aralkylthio, aryloxyalkylthio, arylaminoalkylthio, arylthioalkylthio, alkylamino, cycloalkyl and cycloalkylalkyl;

wherein X₄ and X₇ are independently selected from the group comprising consisting of hydrogen, oxygen, halogen, oxo, carbonyl, thiocarbonyl, hydroxyl, alkyl, aryl, Het¹, Het¹alkyl, Het¹aryl, alkenyl, alkynyl, hydroxyalkyl, hydroxycarbonyl, hydroxycarbonylalkyl, hydroxycarbonylaryl, hydroxycarbonyloxyalkyl, glycosyl, thio derivatives thereof, amino derivatives thereof, carboxy derivatives thereof, amido derivatives thereof, hydroxyl-protected derivatives thereof, optionally substituted by one or more substituents independently selected from the group comprising consisting of alkyl, aralkyl, aryl, Het¹, Het², cycloalkyl, alkyloxy, alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(alkyl)aminocarbonyl, aminosulfonyl, alkylS(=O)_t, hydroxy, cyano, halogen or amino optionally mono- or disubstituted wherein the substituents are independently selected from the group comprising consisting of alkyl, aryl, aralkyl, aryloxy, arylamino, arylthio, aryloxyalkyl, arylaminoalkyl, aralkoxy, alkylthio, alkoxy, aryloxyalkoxy, ~~arylaminealkoxy~~ arylaminoalkoxy, aralkylamino, aryloxyalkylamino, arylaminoalkylamino, arylthioalkoxy, arylthioalkylamino, aralkylthio, aryloxyalkylthio, arylaminoalkylthio, arylthioalkylthio, alkylamino, cycloalkyl and cycloalkylalkyl;

wherein at least one of X₃, X'₃, X₄ and X₇ is a glycosyl moiety; or a deoxy derivative thereof, a carboxy derivative thereof, a hydroxy protected derivative thereof, an amino derivative thereof, an amido derivatives thereof, a thio derivative thereof, optionally substituted by one or more substituents,

wherein X_5 participates ~~to~~in a double bond between the carbon atoms in position 4 and 5 or between carbon atoms in position 5 and 6, and X_6 is selected from the group ~~emprising~~consisting of hydrogen, hydroxyl and hydroxyalkyl, or

wherein X_5 and X_6 are independently selected from the group ~~emprising~~consisting of halogen, hydrogen, hydroxyl, hydroxyalkyl, aminoalkyl, aminoaryl, optionally substituted by one or more substituents independently selected from the group ~~emprising~~consisting of alkyl, aralkyl, aryl, Het¹, Het², cycloalkyl, alkyloxy, alkyloxycarbonyl, carboxyl, aminocarbonyl, and

wherein n is an integer between 0 and 10.

2. (Currently amended) A—The compound according to claim 1, stereoisomers, tautomers, racemics, prodrugs, metabolites thereof, or a pharmaceutically acceptable salt and/or solvate thereof,

wherein X_1 , X_2 , R_1 and R_2 are independently selected from the group ~~emprising~~consisting of oxo, hydrogen, hydroxyl, oxyalkyl, alkyl, alkenyl, alkynyl, alkyloxy, alkyloxyalkyl, alkylthioalkyl, alkoxycarbonyl, alkylthiocarbonyl, alkanoyl, cycloalkylalkyl, cycloalkylcarbonyl, cycloalkylalkanoyl, cycloalkylthiocarbonyl, cycloalkylalkoxycarbonyl, cycloalkylalkoxythiocarbonyl, cycloalkylthioalkyl, alkylcarbonyloxyalkyl, cycloalkylcarbonyloxyalkyl, silyloxyalkyl, aralkyl, arylalkenyl, arylcarbonyl, aryloxycarbonyl, arylthiocarbonyl, aralkoxycarbonyl, arylalkylthiocarbonyl, aryloxyalkyl, arylthioalkyl, haloalkyl, hydroxyalkyl, aralkanoyl, aroyl, aryloxycarbonylalkyl, aryloxyalkanoyl, carboxyl, alkenylcarbonyl, alkynylcarbonyl, Het¹, Het¹alkyl, Het¹oxyalkyl, Het¹aryl, Het¹aralkyl, Het¹cycloalkyl, Het¹alkoxycarbonyl, Het¹alkylthiocarbonyl, Het¹oxycarbonyl, Het¹thiocarbonyl, Het¹alkanoyl, Het¹aralkanoyl, Het¹aryloxyalkyl, Het¹alkyloxyalkyl, Het¹arylthioalkyl, Het¹aryloxycarbonyl, Het¹aralkoxycarbonyl, Het¹aroyl, Het¹oxyalkylcarbonyl, Het¹alkyloxyalkylcarbonyl, Het¹aryloxyalkylcarbonyl, Het¹carbonyloxyalkyl, Het¹alkylcarbonyloxyalkyl, Het¹aralkylcarbonyloxyalkyl, Het²alkyl, Het²oxyalkyl, Het²alkyloxyalkyl, Het²aralkyl, Het²carbonyl, Het²oxycarbonyl, Het²thiocarbonyl, Het²alkanoyl, Het²alkylthiocarbonyl, Het²alkoxycarbonyl, Het²aralkanoyl, Het²aralkoxycarbonyl, Het²aryloxycarbonyl,

Het²aroyl, Het²aryloxyalkyl, Het²arylthioalkyl, Het²oxyalkylcarbonyl, Het²alkyloxyalkylcarbonyl, Het²aryloxyalkylcarbonyl, Het²carbonyloxyalkyl, Het²alkylcarbonyloxyalkyl, Het²aralkylcarbonyloxyalkyl, cyano, CR³=NR⁴, CR³=N(OR⁴), aminocarbonyl, aminoalkanoyl, aminoalkyl, optionally substituted by one or more substituents independently selected from the group ~~comprising~~consisting of alkyl, aralkyl, aryl, Het¹, Het², cycloalkyl, alkyloxy, alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(alkyl)aminocarbonyl, aminosulfonyl, alkylS(=O)_t, hydroxy, cyano, halogen or amino optionally mono- or disubstituted wherein the substituents are independently selected from the group ~~comprising~~consisting of alkyl, aryl, aralkyl, aryloxy, arylamino, arylthio, aryloxyalkyl, arylaminoalkyl, aralkoxy, alkylthio, alkoxy, aryloxyalkoxy, ~~arylaminealkoxy~~arylaminealkoxy, aralkylamino, aryloxyalkylamino, arylaminoalkylamino, arylthioalkoxy, arylthioalkylamino, aralkylthio, aryloxyalkylthio, arylaminoalkylthio, arylthioalkylthio, alkylamino, cycloalkyl, cycloalkylalkyl, Het¹, Het², Het¹alkyl, Het²alkyl, Het¹amino, Het²amino, Het¹alkylamino, Het²alkylamino, Het¹thio, Het²thio, Het¹alkylthio, Het²alkylthio, Het¹oxy and Het²oxy, OR³, SR³, SO₂NR³R⁴, SO₂N(OH)R³, CN, CR³=NR⁴, S(O)R³, SO₂R³, CR³=N(OR⁴), N₃, NO₂, NR³R⁴, N(OH)R³, C(O)R³, C(S)R³, CO₂R³, C(O)SR³, C(O)NR³R⁴, C(S)NR³R⁴, C(O)N(OH)R⁴, C(S)N(OH)R³, NR³C(O)R⁴, NR³C(S)R⁴, N(OH)C(O)R⁴, N(OH)C(S)R³, NR³CO₂R⁴, NR³C(O)NR⁴R⁵, and NR³C(S)NR⁴R⁵, N(OH)CO₂R³, NR³C(O)SR⁴, N(OH)C(O)NR³R⁴, N(OH)C(S)NR³R⁴, NR³C(O)N(OH)R⁴, NR³C(S)N(OH)R⁴, NR³SO₂R⁴, NHSO₂NR³R⁴, NR³SO₂NHR⁴, P(O)(OR³)(OR⁴), wherein t is an integer between 1 and 2 and R³, R⁴ and R⁵ are each independently selected from the group ~~comprising~~consisting of hydrogen, hydroxyl, alkyl, alkenyl, alkynyl, aminoalkyl, aminoaryl, alkylcarbonylamino, arylcarbonylamino, alkylthiocarbonylamino and arylthiocarbonylamino;

wherein X₃ participates together with X'₃ ~~to in~~ an oxo functional group, or wherein X₃ and X'₃ are independently selected from the group ~~comprising~~consisting of hydrogen, hydroxyl, sulfur, oxyalkyl, oxycarbonyl, alkyl, Het¹alkyl, alkyloxycarbonyl, alkenyl, alkynyl, aminoalkyl, aminoacyl, alkylcarbonylamino, alkylthiocarbonylamino, Het¹, glucosyl, fructosyl, galactosyl, mannosyl, ribosyl, ribulosyl, xylulosyl, erythrosyl, erythrulosyl, rhamnosyl, threosyl, sorbosyl, psicosyl, tagatosyl, fucosyl, arabinosyl,

xylofuranosyl, lyxosyl, talosyl, psicose, idosyl, gulose, altrose, allose, mannoheptulosyl, sedoheptulosyl, abequosyl, isomaltosyl, kojibiosyl, laminaribiosyl, nigerosyl, primeverosyl, rutinose, tyvelosyl, maltosyl, lactosyl, sucrose, cellobiosyl, trehalosyl, gentiobiosyl, melibiosyl, turanosyl, sophorose, isosucrose, raffinose, palatinose, lactulosyl, gentianose, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, isomaltotriosyl, maltopentaosyl, maltohexaosyl, maltoheptaosyl, sicosyl, panosyl, isopanosyl, inosyl, N-acetyl-galactosaminyl, mannotriosyl, globotriosyl, erlose, neotrehalose, chitobiosyl, chitobiosemannosyl, glucosaminyl, N-acetyl-glucosaminyl, octylglucopyranosyl, octylribofuranosyl, cyclohexylglucopyranosyl, cyclohexylxylofuranosyl, benzylglucopyranosyl, benzylarabinofuranosyl, N-acetyl-lactosaminyl, acosaminyl, amicetosyl, amylose, apiose, arcanose, ascarylose, bacillosaminyl, boivinosyl, cellotriosyl, chacotriosyl, chalcose, cladinosyl, colitosyl, cymarose, daunosaminyl, desosaminyl, D-glycero-L-gulo-heptosyl, diginose, digitalose, digitoxosyl, evalose, evernitrosyl, forosaminyl, fucosaminyl, garosaminyl, hamamelosyl, isolevoglucosenonyl, kanosaminyl, kansosaminyl, lactosaminyl, lactosediaminyl, fucitolyl, maltulosyl, mannosaminyl, melezitose, mycaminosyl, mycarose, mycinosyl, mycosaminyl, noviosyl, oleandrosyl, paratose, perosaminyl, planteosyl, pneumosaminyl, purpurosaminyl, quinovosaminyl, quinovose, rhamnitolyl, rhamnosaminyl, rhodinosyl, rhodosaminyl, sarmentosyl, solatriosyl, stachyosyl, streptosyl, umbelliferosyl, trehalosaminyl, 1,6-anhydro-D-glucopyranosyl, 1-hydroxy- α -D-allopyranosyl, 2,3:5,6-di-O-isopropylidene-D-mannofuranosyl, 2-amino-2-deoxy-D-galactitolyl, 2-deoxyribose, 2-deoxyglucose, 5-amino-5-deoxy-D-glucopyranosyl, 6-deoxy-D-galactitolyl, 2-amino-2-deoxy glucose, 2-acetamido-2-deoxy-glucose, 2-amino-2-deoxy galactose, 2-acetamido-2-deoxy-galactose, 2-amino-2-deoxy mannose, 2-acetamido-2-deoxy-mannose, 2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucose, 2-amino-2-deoxy-4-O- β -D-galactosyl-D-glucose, 6'-N-acetylglucosaminyl-lactose, 2-acetamido-2-deoxy-3-O- α -L-fucose-D-glucose, 6-O(2-acetamido-2-deoxy- β -D-glucose)-D-galactose, 2-acetamido-2-deoxy-3-O- β -D-galactosyl-D-glucose, 2'-acetamido-2'-deoxy-3-O- β -D-glucose-D-galactose, 3-fucose-D-lactose, 3-fucose-2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucose, L or D isomers thereof, α or β form thereof, pyranuronic or

furanuronic form ~~thereof~~thereof, pyranose or furanose form thereof, combination thereof, deoxy derivatives thereof, hydroxyl-protected acetate or benzoyl derivatives thereof, amino derivatives thereof, amido derivatives thereof, thio derivatives thereof, di-, tri-, oligo- and polysaccharide thereof optionally substituted by one or more substituents independently selected from the group comprisingconsisting of alkyl, aralkyl, aryl, Het¹, Het², cycloalkyl, alkyloxy, alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(alkyl)aminocarbonyl, aminosulfonyl, alkylS(=O)_t, hydroxy, cyano, halogen or amino optionally mono- or disubstituted wherein the substituents are independently selected from the group comprisingconsisting of alkyl, aryl, aralkyl, aryloxy, arylamino, arylthio, aryloxyalkyl, arylaminoalkyl, aralkoxy, alkylthio, alkoxy, aryloxyalkoxy, ~~arylaminoalkoxy~~arylaminoalkoxy, aralkylamino, aryloxyalkylamino, arylaminoalkylamino, arylthioalkoxy, arylthioalkylamino, aralkylthio, aryloxyalkylthio, arylaminoalkylthio, arylthioalkylthio, alkylamino, cycloalkyl and cycloalkylalkyl;

wherein X₄ and X₇ are independently selected from the group comprisingconsisting of hydrogen, oxygen, halogen, oxo, carbonyl, thiocarbonyl, hydroxyl, alkyl, aryl, Het¹, Het¹alkyl, Het¹aryl, alkenyl, alkynyl, hydroxyalkyl, hydroxycarbonyl, hydroxycarbonylalkyl, hydroxycarbonylaryl, hydroxycarbonyloxyalkyl, glucosyl, fructosyl, galactosyl, mannosyl, ribosyl, ribulosyl, xylulosyl, erythrosyl, erythrulosyl, rhamnosyl, threosyl, sorbosyl, psiceryl, tagatosyl, fucosyl, arabinosyl, xylofuranosyl, lyxosyl, talosyl, psiceryl, idosyl, gulosyl, altrosyl, allosyl, mannoheptulosyl, sedoheptulosyl, abequosyl, isomaltosyl, kojibiosyl, laminaribiosyl, nigerosyl, primeverosyl, rutinyl, tyvelosyl, maltosyl, lactosyl, sucrosyl, cellobiosyl, trehalosyl, gentiobiosyl, melibiosyl, turanosyl, sophorosyl, isosucrosyl, raffinyl, palatinosyl, lactulosyl, gentianosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, isomaltotriosyl, maltopentaosyl, maltohexaosyl, maltoheptaosyl, sicosyl, panosyl, isopanosyl, inosyl, N-acetyl-galactosaminyl, mannotriosyl, globotriosyl, erlosyl, neotrehalosyl, chitobiosyl, chitobiosemannosyl, glucosaminyl, N-acetyl-glucosaminyl, octylglucopyranosyl, octylribofuranosyl, cyclohexylglucopyranosyl, cyclohexylxylofuranosyl, benzylglucopyranosyl, benzylarabinofuranosyl, N-acetyl-lactosaminyl, acosaminyl,

amicetosyl, amylosyl, apiosyl, arcanosyl, ascarylosyl, bacillosaminyl, boivinosyl, cellotriosyl, chacotriosyl, chalcosyl, cladinosyl, colitosyl, cymarosyl, daunosaminyl, desosaminyl, D-glycero-L-gulo-heptosyl, diginosyl, digitalosyl, digitoxosyl, evalosyl, evernitrosyl, forosaminyl, fucosaminyl, garosaminyl, hamamelosyl, isolevoglucosenonyl, kanosaminyl, kansosaminyl, lactosaminyl, lactosediainyl, fucitolyl, maltulosyl, mannosaminyl, melezitoyl, mycaminosyl, mycarosyl, mycinosyl, mycosaminyl, noviosyl, oleandrosyl, paratosyl, perosaminyl, planteosyl, pneumosaminyl, purpurosaminyl, quinovosaminyl, quinovosyl, rhamnitolyl, rhamnosaminyl, rhodinosyl, rhodosaminyl, sarmentosyl, solatriosyl, stachyosyl, streptosyl, umbelliferosyl, trehalosaminyl, 1,6-anhydro-D-glucopyranosyl, 1-hydroxy- α -D-allopyranosyl, 2,3:5,6-di-O-isopropylidene-D-mannofuranosyl, 2-amino-2-deoxy-D-galactitolyl, 2-deoxyribosyl, 2-deoxyglucosyl, 5-amino-5-deoxy-D-glucopyranosyl, 6-deoxy-D-galactitolyl, 2-amino-2-deoxy glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-amino-2-deoxy-mannosyl, 2-acetamido-2-deoxy-mannosyl, 2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminylactosyl, 2-acetamido-2-deoxy-3-O- α -L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy- β -D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O- β -D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O- β -D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form thereof, pyranose or furanose form thereof, combination thereof, deoxy derivatives thereof, hydroxyl-protected acetate or benzoyl derivatives thereof, amino derivatives thereof, amido derivatives thereof, thio derivatives thereof, di-, tri-, oligo- and polysaccharide thereof optionally substituted by one or more substituents independently selected from the group ~~comprising~~consisting of alkyl, aralkyl, aryl, Het¹, Het², cycloalkyl, alkyloxy, alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(alkyl)aminocarbonyl, aminosulfonyl, alkylS(=O)_t, hydroxy, cyano, halogen or amino optionally mono- or disubstituted wherein the substituents are independently selected from the group ~~comprising~~consisting of alkyl, aryl, aralkyl, aryloxy, arylamino, arylthio, aryloxyalkyl, arylaminoalkyl, aralkoxy, alkylthio, alkoxy, aryloxyalkoxy, ~~arylaminealkoxy~~arylaminealkoxy, aralkylamino, aryloxyalkylamino, arylaminealkylamino,

arylthioalkoxy, arylthioalkylamino, aralkylthio, aryloxyalkylthio, arylaminoalkylthio, arylthioalkylthio, alkylamino, cycloalkyl and cycloalkylalkyl;

wherein at least one of X_3 , X'_3 , X_4 and X_7 is a glycosyl moiety selected from the group ~~comprising~~consisting of glucosyl, fructosyl, galactosyl, mannosyl, ribosyl, ribulosyl, xylulosyl, erythrosyl, erythrulosyl, rhamnosyl, threosyl, sorbosyl, psicose, tagatosyl, fucosyl, arabinosyl, xylofuranosyl, lyxosyl, talosyl, psicose, idosyl, gulonic, allosyl, mannoheptulosyl, sedoheptulosyl, abequosyl, isomaltosyl, kojibiosyl, laminaribiosyl, nigerosyl, primeverosyl, rutinyl, tyvelosyl, maltosyl, lactosyl, sucrosyl, cellobiosyl, trehalosyl, gentiobiosyl, melibiosyl, turanosyl, sophorosyl, isosucrosyl, raffinyl, palatinosyl, lactulosyl, gentianosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, isomaltotriosyl, maltopentaosyl, maltohexaosyl, maltoheptaosyl, sicosyl, panosyl, isopanosyl, inosyl, N-acetyl-galactosaminyl, mannotriosyl, globotriosyl, erlosyl, neotrehalosyl, chitobiosyl, chitobiosemannosyl, glucosaminyl, N-acetyl-glucosaminyl, octylglucopyranosyl, octylribofuranosyl, cyclohexylglucopyranosyl, cyclohexylxylofuranosyl, benzylglucopyranosyl, benzylarabinofuranosyl, N-acetyl-lactosaminyl, acosaminyl, amicetosyl, amylosyl, apiosyl, arcanosyl, ascarylosyl, bacillosaminyl, boivinosyl, cellosyl, chacotriosyl, chalcosyl, cladinosyl, colitosyl, cymarosyl, daunosaminyl, desosaminyl, D-glycero-L-gulo-heptosyl, diginosyl, digitalosyl, digitoxosyl, evalosyl, evernitrosyl, forosaminyl, fucosaminyl, garosaminyl, hamamelosyl, isolevoglucosenonyl, kanosaminyl, kansosaminyl, lactosaminyl, lactosediaminyl, fucitolyl, maltulosyl, mannosaminyl, melezitoyl, mycaminosyl, mycarosyl, mycinosyl, mycosaminyl, noviosyl, oleandrosyl, paratosyl, perosaminyl, planteosyl, pneumosaminyl, purpurosaminyl, quinovosaminyl, quinovosyl, rhamnitolyl, rhamnosaminyl, rhodinosyl, rhodosaminyl, sarmentosyl, solatriosyl, stachyosyl, streptosyl, umbelliferosyl, trehalosaminyl, 1,6-anhydro-D-glucopyranosyl, 1-hydroxy- α -D-allopyranosyl, 2,3:5,6-di-O-isopropylidene-D-mannofuranosyl, 2-amino-2-deoxy-D-galactitolyl, 2-deoxyribosyl, 2-deoxyglucosyl, 5-amino-5-deoxy-D-glucopyranosyl, 6-deoxy-D-galactitolyl, 2-amino-2-deoxy glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-amino-2-deoxy mannosyl, 2-acetamido-2-deoxy-

mannosyl, 2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminylactosyl, 2-acetamido-2-deoxy-3-O- α -L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy- β -D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O- β -D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O- β -D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form thereof, pyranose or furanose form thereof, combination thereof, deoxy derivatives thereof, hydroxyl-protected acetate or benzoyl derivatives thereof, amino derivatives thereof, amido derivatives thereof, thio derivatives thereof, di-, tri-, oligo- and polysaccharide thereof optionally substituted by one or more substituents independently selected from the group consisting of alkyl, aralkyl, aryl, Het¹, Het², cycloalkyl, alkyloxy, alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(alkyl)aminocarbonyl, aminosulfonyl, alkylS(=O)_t, hydroxy, cyano, halogen or amino optionally mono- or disubstituted wherein the substituents are independently selected from the group consisting of alkyl, aryl, aralkyl, aryloxy, arylamino, arylthio, aryloxyalkyl, arylaminoalkyl, aralkoxy, alkylthio, alkoxy, aryloxyalkoxy, arylaminoalkoxy, aralkylamino, aryloxyalkylamino, arylaminoalkylamino, arylthioalkoxy, arylthioalkylamino, aralkylthio, aryloxyalkylthio, arylaminoalkylthio, arylthioalkylthio, alkylamino, cycloalkyl and cycloalkylalkylas indicated above ;

wherein X₅ participates to in a double bond between the carbon atoms in position 4 and 5 or between carbon atoms in positions 5 and 6, and X₆ is selected from the group comprising consisting of hydrogen, hydroxyl and hydroxyalkyl, or

wherein X₅ and X₆ are independently selected from the group comprising consisting of halogen, hydrogen, hydroxyl, hydroxyalkyl, aminoalkyl, aminoaryl, optionally substituted by one or more substituents independently selected from the group comprising consisting of alkyl, aralkyl, aryl, Het¹, Het², cycloalkyl, alkyloxy, alkyloxycarbonyl, carboxyl, aminocarbonyl, and

wherein n is an integer between 0 and 10.

3. (Currently amended) A The compound according to claim 1[[or 2]], stereoisomers, tautomers, racemics, prodrugs, metabolites thereof, or a pharmaceutically acceptable salt and/or solvate thereof,

wherein X_1 , X_2 , R_1 and R_2 are independently selected from the group comprising~~consisting of~~ hydrogen, hydroxyl, oxyalkyl, oxo, alkyl, alkenyl, alkynyl, alkyloxy, alkyloxyalkyl, alkylthioalkyl, alkoxycarbonyl, alkylthiocarbonyl, alkanoyl, cycloalkylalkyl, cycloalkylcarbonyl, cycloalkylalkanoyl, cycloalkylthiocarbonyl, cycloalkylalkoxycarbonyl, cycloalkylalkoxythiocarbonyl, cycloalkylthioalkyl, alkylcarbonyloxyalkyl, cycloalkylcarbonyloxyalkyl, silyloxyalkyl, aralkyl, arylalkenyl, arylcarbonyl, aryloxycarbonyl, arylthiocarbonyl, aralkoxycarbonyl, arylalkylthiocarbonyl, aryloxyalkyl, arylthioalkyl, haloalkyl, hydroxyalkyl, aralkanoyl, aroyl, aryloxycarbonylalkyl, aryloxyalkanoyl, carboxyl, alkenylcarbonyl and alkynylcarbonyl;

wherein X_3 participates together with X'_3 ~~to~~in an oxo functional group, or wherein X_3 and X'_3 are independently selected from the group comprising~~consisting of~~ hydrogen, hydroxyl, sulfur, oxyalkyl, oxycarbonyl, alkyl, Het¹alkyl, alkyloxycarbonyl, alkenyl, alkynyl, aminoalkyl, aminoacyl, alkylcarbonylamino, alkylthiocarbonylamino, Het¹, glucosyl, fructosyl, galactosyl, mannosyl, ribosyl, ribulosyl, xylulosyl, erythrosyl, erythrosyl, rhamnosyl, threosyl, sorbosyl, psicosyl, tagatosyl, fucosyl, arabinosyl, xylofuranosyl, lyxosyl, talosyl, psicosyl, idosyl, gulosyl, altrosyl, allosyl, mannoheptulosyl, sedoheptulosyl, abequosyl, isomaltosyl, kojibiosyl, laminaribiosyl, nigerosyl, primeverosyl, rutiniosyl, tyvelosyl, maltosyl, lactosyl, sucrosyl, cellobiosyl, trehalosyl, gentiobiosyl, melibiosyl, turanosyl, sophorosyl, isosucrosyl, raffiniosyl, palatinosyl, lactulosyl, gentianiosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, 2-amino-2-deoxy glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy-galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-amino-2-deoxy-mannosyl, 2-acetamido-2-deoxy-mannosyl, 2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminyl lactosyl, 2-acetamido-2-deoxy-3-O- α -L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy- β -D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O- β -D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O- β -D-glucosyl-D-galactosyl, 3-fucosyl-

D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form thereof, pyranose or furanose form thereof, combination thereof, deoxy derivatives thereof, hydroxyl-protected acetate or benzoyl derivatives thereof, amino derivatives thereof, amido derivatives thereof, thio derivatives thereof, disaccharide thereof, trisaccharide thereof, oligosaccharide and polysaccharide thereof optionally substituted by one or more substituents independently selected from the group ~~comprising~~consisting of alkyl, aralkyl, aryl, Het¹, Het², cycloalkyl, alkyloxy, alkyloxycarbonyl, carboxyl and aminocarbonyl;

wherein X₄ and X₇ are independently selected from the group ~~comprising~~consisting of hydrogen, oxygen, halogen, oxo, carbonyl, thiocarbonyl, hydroxyl, alkyl, aryl, Het¹, Het¹alkyl, Het¹aryl, alkenyl, alkynyl, hydroxyalkyl, hydroxycarbonyl, hydroxycarbonylalkyl, hydroxycarbonylaryl, hydroxycarbonyloxyalkyl, glucosyl, fructosyl, galactosyl, mannosyl, ribosyl, ribulosyl, xylulosyl, erythrosyl, erythrulosyl, rhamnosyl, threosyl, sorbosyl, psiceryl, tagatosyl, fucosyl, arabinosyl, xylofuranosyl, lyxosyl, talosyl, psiceryl, idosyl, gulosyl, altrosyl, allosyl, mannoheptulosyl, sedoheptulosyl, abequosyl, isomaltosyl, kojibiosyl, laminaribiosyl, nigerosyl, primeverosyl, rutinyl, tyvelosyl, maltosyl, lactosyl, sucrosyl, cellobiosyl, trehalosyl, gentiobiosyl, melibiosyl, turanosyl, sophorosyl, isosucrosyl, raffinyl, palatinosyl, lactulosyl, gentianosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, 2-amino-2-deoxy-glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-amino-2-deoxy-mannosyl, 2-acetamido-2-deoxy-mannosyl, 2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminyl lactosyl, 2-acetamido-2-deoxy-3-O- α -L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy- β -D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O- β -D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O- β -D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form thereof, pyranose or furanose form thereof, combination thereof, deoxy derivatives thereof, hydroxyl-protected acetate or benzoyl derivatives thereof, amino derivatives thereof,

amido derivatives thereof, thio derivatives thereof, disaccharide thereof, trisaccharide thereof, oligosaccharide and polysaccharide thereof optionally substituted by one or more substituents independently selected from the group comprising consisting of alkyl, aralkyl, aryl, Het¹, Het², cycloalkyl, alkyloxy, alkyloxycarbonyl, carboxyl and aminocarbonyl;

wherein at least one of X₃, X'₃, X₄ and X₇ is a glycosyl moiety selected from the group- consisting of glucosyl, fructosyl, galactosyl, mannosyl, ribosyl, ribulosyl, xylulosyl, erythrosyl, erythrulosyl, rhamnosyl, threosyl, sorbosyl, psicose, tagatosyl, fucosyl, arabinosyl, xylofuranosyl, lyxosyl, talosyl, psicose, idosyl, gulose, altrose, allose, mannoheptulosyl, sedoheptulosyl, abequosyl, isomaltosyl, kojibiosyl, laminaribiosyl, nigerosyl, primeverosyl, rutinose, tyvelosyl, maltosyl, lactosyl, sucrose, cellobiosyl, trehalosyl, gentiobiosyl, melibiosyl, turanose, sophorose, isosucrose, raffinose, palatinose, lactulose, gentianose, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, isomaltotriosyl, maltopentaosyl, maltohexaosyl, maltoheptaosyl, sicosyl, panosyl, isopanose, inosyl, N-acetyl-galactosaminyl, mannotriosyl, globotriosyl, erlose, neotrehalose, chitobiosyl, chitobiosemannosyl, glucosaminyl, N-acetyl-glucosaminyl, octylglucopyranosyl, octylribofuranosyl, cyclohexylglucopyranosyl, cyclohexylxylofuranosyl, benzylglucopyranosyl, benzylarabinofuranosyl, N-acetyl-lactosaminyl, acosaminyl, amicetose, amylose, apiose, arcanose, ascarylose, bacillosaminyl, bovinosyl, cellotriosyl, chacotriosyl, chalcose, cladinosyl, colitose, cymarose, daunosaminyl, desosaminyl, D-glycero-L-gulo-heptosyl, diginose, digitalose, digitoxose, evalose, evernitrosyl, forosaminyl, fucosaminyl, garosaminyl, hamamelose, isolevoglucosenonyl, kanosaminyl, kansosaminyl, lactosaminyl, lactosedi-aminyl, fucitolyl, maltulose, mannosaminyl, melezitose, mycaminosyl, mycarose, mycinose, mycosaminyl, noviose, oleandrose, paratose, perosaminyl, planteose, pneumosaminyl, purpurosaminyl, quinosaminyl, quinosyl, rhamnitolyl, rhamnosaminyl, rhodinosyl, rhodosaminyl, sarmentose, solatriosyl, stachyose, streptose, umbelliferose, trehalosaminyl, 1,6-anhydro-D-glucopyranosyl, 1-hydroxy- α -D-allopyranosyl, 2,3:5,6-di-O-isopropylidene-D-mannofuranosyl, 2-amino-2-deoxy-D-galactitolyl, 2-deoxyribosyl, 2-deoxyglucose, 5-amino-5-deoxy-D-glucopyranosyl, 6-deoxy-D-galactitolyl, 2-amino-2-

deoxy glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-amino-2-deoxy mannosyl, 2-acetamido-2-deoxy-mannosyl, 2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminylactosyl, 2-acetamido-2-deoxy-3-O- α -L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy- β -D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O- β -D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O- β -D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form thereof, pyranose or furanose form thereof, combination thereof, deoxy derivatives thereof, hydroxyl-protected acetate or benzoyl derivatives thereof, amino derivatives thereof, amido derivatives thereof, thio derivatives thereof, di-, tri-, oligo- and polysaccharide thereof optionally substituted by one or more substituents independently selected from the group consisting of alkyl, aralkyl, aryl, Het¹, Het², cycloalkyl, alkyloxy, alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(alkyl)aminocarbonyl, aminosulfonyl, alkylS(=O)_t, hydroxy, cyano, halogen or amino optionally mono- or disubstituted wherein the substituents are independently selected from the group consisting of alkyl, aryl, aralkyl, aryloxy, arylamino, arylthio, aryloxyalkyl, arylaminoalkyl, aralkoxy, alkylthio, alkoxy, aryloxyalkoxy, arylaminoalkoxy, aralkylamino, aryloxyalkylamino, arylaminoalkylamino, arylthioalkoxy, arylthioalkylamino, aralkylthio, aryloxyalkylthio, arylaminoalkylthio, arylthioalkylthio, alkylamino, cycloalkyl and cycloalkylalkyl as indicated above;

wherein X₅ participates to ~~in~~ a double bond between the carbon atoms in position 4 and 5 or between carbon atoms in position 5 and 6, and X₆ is selected from the group ~~comprising~~ consisting of hydrogen, hydroxyl, and hydroxyalkyl, or wherein X₅ and X₆ are independently selected from the group ~~comprising~~ consisting of hydrogen, hydroxyl, hydroxyalkyl, aminoalkyl, aminoaryl, optionally substituted by one or more substituents independently selected from the group ~~comprising~~ consisting of alkyl, aralkyl, aryl, Het¹, Het², cycloalkyl, alkyloxy, alkyloxycarbonyl, carboxyl, aminocarbonyl, and

wherein n is an integer between 0 and 5.

4. (Currently amended) A ~~The~~ compound according to ~~any of claims 1 to 3~~ claim 1, stereoisomers, tautomers, racemics, prodrugs, metabolites thereof, or a pharmaceutically acceptable salt and/or solvate thereof,

wherein X_1 , X_2 , R_1 and R_2 are independently selected from the group ~~comprising~~ consisting of hydrogen, hydroxyl, alkyloxy, oxo and oxyalkyl,

wherein X_3 participates together with X'_3 ~~to in~~ in an oxo functional group, or wherein X_3 and X'_3 are independently selected from the group ~~comprising~~ consisting of hydrogen, hydroxyl, oxyalkyl, oxycarbonyl, glucosyl, fructosyl, galactosyl, mannosyl, ribosyl, ribulosyl, xylulosyl, erythrosyl, erythrulosyl, rhamnosyl, threosyl, sorbosyl, psicose, tagatose, fucose, arabinose, altrose, laminaribiosyl, isomaltosyl, maltosyl, lactosyl, cellobiosyl, gentiobiosyl, melibiosyl, palatinosyl, lactulosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, 2-amino-2-deoxy-glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy-galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-amino-2-deoxy-mannosyl, 2-acetamido-2-deoxy-mannosyl, 2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminyl-lactosyl, 2-acetamido-2-deoxy-3-O- α -L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy- β -D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O- β -D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O- β -D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form ~~thereof~~ thereof, pyranose or furanose form thereof, combination thereof, deoxy derivatives thereof, hydroxyl-protected acetate derivatives thereof, amino derivatives thereof, amido derivatives thereof, thio derivatives thereof, disaccharide thereof, trisaccharide thereof, oligosaccharide and polysaccharide thereof;

wherein X_4 and X_7 are independently selected from the group ~~comprising~~ consisting of hydrogen, oxygen, oxo, hydroxyl, glucosyl, fructosyl, galactosyl, mannosyl, ribosyl, ribulosyl, xylulosyl, erythrosyl, erythrulosyl, rhamnosyl, threosyl, sorbosyl, psicose, tagatose, fucose, arabinose, altrose, laminaribiosyl, isomaltosyl, maltosyl, lactosyl, cellobiosyl, gentiobiosyl, melibiosyl, palatinosyl, lactulosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl,

maltotetraosyl, 2-amino-2-deoxy-glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-amino-2-deoxy-mannosyl, 2-acetamido-2-deoxy-mannosyl, 2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminylactosyl, 2-acetamido-2-deoxy-3-O- α -L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy- β -D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O- β -D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O- β -D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form thereof, pyranose or furanose form thereof, combination thereof, deoxy derivatives thereof, hydroxyl-protected acetate or benzoyl derivatives thereof, amino derivatives thereof, amido derivatives thereof, thio derivatives thereof, disaccharide thereof, trisaccharide thereof, oligosaccharide and polysaccharide thereof;

wherein at least one of X_3 , X'_3 , X_4 and X_7 is a glycosyl moiety selected from the group consisting of glucosyl, fructosyl, galactosyl, mannosyl, ribosyl, ribulosyl, xylulosyl, erythrosyl, erythrulosyl, rhamnosyl, threosyl, sorbosyl, psiceryl, tagatosyl, fucosyl, arabinosyl, xylofuranosyl, lyxosyl, talosyl, psiceryl, idosyl, gulosyl, altrosyl, allosyl, mannoheptulosyl, sedoheptulosyl, abequosyl, isomaltosyl, kojibiosyl, laminaribiosyl, nigerosyl, primeverosyl, rutinolyl, tyvelosyl, maltosyl, lactosyl, sucrosyl, cellobiosyl, trehalosyl, gentiobiosyl, melibiosyl, turanosyl, sophorosyl, isosucrosyl, raffinolyl, palatinosyl, lactulosyl, gentianolyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, isomaltotriosyl, maltopentaosyl, maltohexaosyl, maltoheptaosyl, sicosyl, panosyl, isopanosyl, inosyl, N-acetylglactosaminyl, mannotriosyl, globotriosyl, erlosyl, neotrehalosyl, chitobiosyl, chitobiosemannosyl, glucosaminyl, N-acetyl-glucosaminyl, octylglucopyranosyl, octylribofuranosyl, cyclohexylglucopyranosyl, cyclohexylxylofuranosyl, benzylglucopyranosyl, benzylarabinofuranosyl, N-acetyl-lactosaminyl, acosaminyl, amicetosyl, amylosyl, apiosyl, arcanosyl, ascarylosyl, bacillosaminyl, boivinosyl, cellotriosyl, chacotriosyl, chalcosyl, cladinosyl, colitosyl, cymarosyl, daunosaminyl, desosaminyl, D-glycero-L-gulo-heptosyl, diginosyl, digitalosyl, digitoxosyl, evalosyl, evernitrosyl, forosaminyl, fucosaminyl, garosaminyl, hamamelosyl, isolevoglucosenonyl,

kanosaminy, kansosaminy, lactosaminy, lactosediaminy, fucitoly, maltulosy,
mannosaminy, melezitoy, mycaminosy, mycarosy, mycinosy, mycosaminy,
noviosy, oleandrosy, paratosy, perosaminy, planteosy, pneumosaminy,
purpurosaminy, quinovosaminy, quinovosy, rhamnity, rhamnosaminy, rhodinosy,
rhodosaminy, sarmentosy, solatriosy, stachyosy, streptosy, umbelliferosy,
trehalosaminy, 1,6-anhydro-D-glucopyranosy, 1-hydroxy- α -D-allopyranosy, 2,3:5,6-di-
O-isopropylidene-D-mannofuranosy, 2-amino-2-deoxy-D-galactity, 2-deoxyribosy, 2-
deoxyglucosy, 5-amino-5-deoxy-D-glucopyranosy, 6-deoxy-D-galactity, 2-amino-2-
deoxy glucosy, 2-acetamido-2-deoxy-glucosy, 2-amino-2-deoxy galactosy, 2-
acetamido-2-deoxy-galactosy, 2-amino-2-deoxy mannosy, 2-acetamido-2-deoxy-
mannosy, 2-acetamido-2-deoxy-4-O- β -D-galactosy-D-glucosy, 2-amino-2-deoxy-4-O-
 β -D-galactosy-D-glucosy, 6'-N-acetylglucosaminyllactosy, 2-acetamido-2-deoxy-3-O-
 α -L-fucosy-D-glucosy, 6-O(2-acetamido-2-deoxy- β -D-glucosy)-D-galactosy, 2-
acetamido-2-deoxy-3-O- β -D-galactosy-D-glucosy, 2'-acetamido-2'-deoxy-3-O- β -D-
glucosy-D-galactosy, 3-fucosy-D-lactosy, 3-fucosy-2-acetamido-2-deoxy-4-O- β -D-
galactosy-D-glucosy, L or D isomers thereof, α or β form thereof, pyranuronic or
furanuronic form thereof, pyranose or furanose form thereof, combination thereof,
deoxy derivatives thereof, hydroxyl-protected acetate or benzoyl derivatives thereof,
amino derivatives thereof, amido derivatives thereof, thio derivatives thereof, di-, tri-,
oligo- and polysaccharide thereof optionally substituted by one or more substituents
independently selected from the group consisting of alkyl, aralkyl, aryl, Het¹, Het²,
cycloalkyl, alkyloxy, alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or
di(alkyl)aminocarbonyl, aminosulfonyl, alkylS(=O)t, hydroxy, cyano, halogen or amino
optionally mono- or disubstituted wherein the substituents are independently selected
from the group consisting of alkyl, aryl, aralkyl, aryloxy, arylamino, arylthio, aryloxyalkyl,
arylaminalkyl, aralkoxy, alkylthio, alkoxy, aryloxyalkoxy, arylaminalkoxy, aralkylamino,
aryloxyalkylamino, arylaminalkylamino, arylthioalkoxy, arylthioalkylamino, aralkylthio,
aryloxyalkylthio, arylaminalkylthio, arylthioalkylthio, alkylamino, cycloalkyl and
cycloalkylalkylas indicated above;

wherein X₄ or X₆ are hydrogen and wherein X₅ participates to in a double bond between the carbon atoms in position 4 and 5 or in position 5 and 6, and

wherein n is an integer between 0 and 2.

5. (Currently amended) A-~~The~~ compound according to ~~any of claims 1 to 4~~claim 1, stereoisomers, tautomers, racemics, prodrugs, metabolites thereof, or a pharmaceutically acceptable salt and/or solvate thereof, wherein X_1 and X_2 are -OMe, wherein R_1 and R_2 are -H, wherein X_3 is selected from the group ~~comprising~~consisting of glucosyl, fructosyl, galactosyl, mannosyl, fucosyl, isomaltosyl, maltosyl, cellobiosyl, gentiobiosyl, melibiosyl, palatinosyl, lactulosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, 2-amino-2-deoxy glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminyllactosyl, 2-acetamido-2-deoxy-3-O- α -L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy- β -D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O- β -D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O- β -D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form ~~thereof~~thereof, pyranose or furanose form thereof, a disaccharide or a trisaccharide thereof, wherein X'_3 is selected from the group ~~comprising~~consisting of hydrogen, alkyl or aralkyl, wherein X_4 is hydrogen, wherein X_5 participates ~~to in~~in a double bond between the carbon atoms in position 5 and 6, wherein X_6 is -H, wherein X_7 is selected from the group ~~comprising~~consisting of hydrogen, oxygen, hydroxyl or oxo, and wherein n is 0.

6. (Currently amended) A-~~The~~ compound according to ~~any of claims 1 to 4~~claim 1, stereoisomers, tautomers, racemics, prodrugs, metabolites thereof, or a pharmaceutically acceptable salt and/or solvate thereof, wherein X_1 and X_2 are -OMe, wherein R_1 and R_2 are -H, wherein X_3 is selected from the group ~~comprising~~consisting of hydrogen, hydroxyl, oxyalkyl or oxycarbonyl, wherein X'_3 is selected from the group ~~comprising~~consisting of glucosyl, fructosyl, galactosyl, mannosyl, fucosyl, isomaltosyl, maltosyl, cellobiosyl, gentiobiosyl, melibiosyl, palatinosyl, lactulosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, 2-amino-2-deoxy-glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy-galactosyl, 2-

acetamido-2-deoxy-galactosyl, 2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 2-Amino-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminylactosyl, 2-acetamido-2-deoxy-3-O- α -L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy- β -D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O- β -D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O- β -D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form ~~thereof~~thereof, pyranose or furanose form thereof, a disaccharide or a trisaccharide thereof, wherein X_4 is hydrogen, wherein X_5 participates ~~to~~in a double bond between the carbon atoms in position 5 and 6, wherein X_6 is -H, wherein X_7 is selected from the group ~~comprising~~consisting of hydrogen, oxygen, hydroxyl or oxo, and wherein n is 0.

7. (Currently amended) A ~~The~~ compound according to ~~any of claims 1 to 4~~claim 1, stereoisomers, tautomers, racemics, prodrugs, metabolites thereof, or a pharmaceutically acceptable salt and/or solvate thereof, wherein X_1 and X_2 are -OMe, wherein R_1 and R_2 are -H, wherein X_3 participates together with X'_3 ~~to~~in an oxo functional group, wherein X_4 is selected from the group ~~comprising~~consisting of glucosyl, fructosyl, galactosyl, mannosyl, fucosyl, isomaltosyl, maltosyl, cellobiosyl, gentiobiosyl, melibiosyl, palatinosyl, lactulosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, 2-amino-2-deoxy-glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy-galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminylactosyl, 2-acetamido-2-deoxy-3-O- α -L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy- β -D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O- β -D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O- β -D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form ~~thereof~~thereof, pyranose or furanose form thereof, a disaccharide or a trisaccharide thereof, wherein X_5 participates ~~to~~in a double bond between the carbon atoms in position 4 and 5, wherein X_6 is -H, wherein X_7 is selected from the group

~~comprising~~consisting of hydrogen, oxygen, hydroxyl, alkyloxy or oxo, and wherein n is 0.

8. (Currently amended) A ~~The~~ compound according to ~~any of claims 1 to 4~~claim 1, stereoisomers, tautomers, racemics, prodrugs, metabolites thereof, or a pharmaceutically acceptable salt and/or solvate thereof, wherein X_1 and X_2 are -OMe, wherein R_1 and R_2 are -H, wherein X_3 participates together with X'_3 ~~to~~in an oxo functional group, wherein X_4 is hydrogen, wherein X_5 participates ~~to~~in a double bond between the carbon atoms in position 5 and 6, wherein X_6 is -H, wherein X_7 is selected from the group ~~comprising~~consisting of glucosyl, fructosyl, galactosyl, mannosyl, fucosyl, isomaltosyl, maltosyl, cellobiosyl, gentiobiosyl, melibiosyl, palatinosyl, lactulosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, 2-amino-2-deoxy glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy-galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminylactosyl, 2-acetamido-2-deoxy-3-O- α -L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy- β -D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O- β -D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O- β -D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form ~~thereof~~thereof, pyranose or furanose form thereof, a disaccharide or a trisaccharide thereof; and wherein n is 0.

9. (Currently amended) A ~~The~~ compound according to ~~any of claims 1 to 4~~claim 1, stereoisomers, tautomers, racemics, prodrugs, metabolites thereof, or a pharmaceutically acceptable salt and/or solvate thereof, wherein X_1 and X_2 are -OMe, wherein R_1 and R_2 are -H, wherein X_3 or X'_3 are independently selected from the group ~~comprising~~consisting of hydrogen or glucosyl, fructosyl, galactosyl, mannosyl, fucosyl, isomaltosyl, maltosyl, cellobiosyl, gentiobiosyl, melibiosyl, palatinosyl, lactulosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, 2-amino-2-deoxy glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-

deoxy-galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 2-Amino-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminylactosyl, 2-acetamido-2-deoxy-3-O- α -L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy- β -D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O- β -D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O- β -D-glucosyl-D-galactosyl, 3'-Fucosyl-D-Lactosyl, 3'-Fucosyl-2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form thereof, pyranose or furanose form thereof, a disaccharide or a trisaccharide thereof, wherein X_4 is selected from the group ~~comprising~~consisting of glucosyl, fructosyl, galactosyl, mannosyl, fucosyl, isomaltosyl, isomaltosyl, cellobiosyl, gentiobiosyl, melibiosyl, palatinosyl, lactulosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, 2-amino-2-deoxy glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy-galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminylactosyl, 2-acetamido-2-deoxy-3-O- α -L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy- β -D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O- β -D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O- β -D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form thereof, pyranose or furanose form thereof, a disaccharide or a trisaccharide thereof, wherein X_5 and X_6 participates to ~~in~~a double bond between the carbon atoms in position 4 and 5, wherein X_6 is -H, wherein X_7 is selected from the group ~~comprising~~consisting of hydrogen, oxygen, hydroxyl, alkyloxy or oxo, wherein at least one of X_3 and X'_3 is a glycosyl moiety selected from the group consisting of glucosyl, fructosyl, galactosyl, mannosyl, ribosyl, ribulosyl, xylulosyl, erythrosyl, erythrulosyl, rhamnosyl, threosyl, sorbosyl, psicosyl, tagatosyl, fucosyl, arabinosyl, xylofuranosyl, lyxosyl, talosyl, psicosyl, idosyl, gulosyl, altrosyl, allosyl, mannoheptulosyl, sedoheptulosyl, abequosyl, isomaltosyl, kojibiosyl, laminaribiosyl, nigerosyl, primeverosyl, rutinoyl, tyvelosyl, maltosyl, lactosyl, sucrosyl, cellobiosyl, trehalosyl, gentiobiosyl, melibiosyl, turanosyl, sophorosyl, isosucrosyl, raffinoyl, palatinosyl, lactulosyl, gentianoyl, 3-mannobiosyl,

6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, isomaltotriosyl, maltopentaosyl, maltohexaosyl, maltoheptaosyl, sicosyl, panosyl, isopanosyl, inosyl, N-acetylgalactosaminyl, mannotriosyl, globotriosyl, erlosyl, neotrehalosyl, chitobiosyl, chitobiosemannosyl, glucosaminyl, N-acetyl-glucosaminyl, octylglucopyranosyl, octylribofuranosyl, cyclohexylglucopyranosyl, cyclohexylxylofuranosyl, benzylglucopyranosyl, benzylarabinofuranosyl, N-acetyl-lactosaminyl, acosaminyl, amicetosyl, amylosyl, apiosyl, arcanosyl, ascarylosyl, bacillosaminyl, boivinosyl, cellotriosyl, chacotriosyl, chalcosyl, cladinosyl, colitosyl, cymarosyl, daunosaminyl, desosaminyl, D-glycero-L-gulo-heptosyl, diginosyl, digitalosyl, digitoxosyl, evalosyl, evernitrosyl, forosaminyl, fucosaminyl, garosaminyl, hamamelosyl, isolevoglucosenonyl, kanosaminyl, kansosaminyl, lactosaminyl, lactosediaminyl, fucitolyl, maltulosyl, mannosaminyl, melezitoyl, mycaminosyl, mycarosyl, mycinosyl, mycosaminyl, noviosyl, oleandrosyl, paratosyl, perosaminyl, planteosyl, pneumosaminyl, purpurosaminyl, quinovosaminyl, quinovosyl, rhamnitolyl, rhamnosaminyl, rhodinosyl, rhodosaminyl, sarmentosyl, solatriosyl, stachyosyl, streptosyl, umbelliferosyl, trehalosaminyl, 1,6-anhydro-D-glucopyranosyl, 1-hydroxy- α -D-allopyranosyl, 2,3:5,6-di-O-isopropylidene-D-mannofuranosyl, 2-amino-2-deoxy-D-galactitolyl, 2-deoxyribosyl, 2-deoxyglucosyl, 5-amino-5-deoxy-D-glucopyranosyl, 6-deoxy-D-galactitolyl, 2-amino-2-deoxy glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-amino-2-deoxy mannosyl, 2-acetamido-2-deoxy-mannosyl, 2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminyl-lactosyl, 2-acetamido-2-deoxy-3-O- α -L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy- β -D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O- β -D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O- β -D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form thereof, pyranose or furanose form thereof, combination thereof, deoxy derivatives thereof, hydroxyl-protected acetate or benzoyl derivatives thereof, amino derivatives thereof, amido derivatives thereof, thio derivatives thereof, di-, tri-, oligo- and polysaccharide thereof optionally substituted by one or more substituents independently selected from the group consisting of alkyl, aralkyl, aryl,

Het¹, Het², cycloalkyl, alkyloxy, alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(alkyl)aminocarbonyl, aminosulfonyl, alkylS(=O)t, hydroxy, cyano, halogen or amino optionally mono- or disubstituted wherein the substituents are independently selected from the group consisting of alkyl, aryl, aralkyl, aryloxy, arylamino, arylthio, aryloxyalkyl, arylaminoalkyl, aralkoxy, alkylthio, alkoxy, aryloxyalkoxy, arylaminoalkoxy, aralkylamino, aryloxyalkylamino, arylaminoalkylamino, arylthioalkoxy, arylthioalkylamino, aralkylthio, aryloxyalkylthio, arylaminoalkylthio, arylthioalkylthio, alkylamino, cycloalkyl and cycloalkylalkylas indicated above and wherein n is 0.

10. (Currently amended) A ~~The~~ compound according to ~~any of claims 1 to 4~~claim 1, stereoisomers, tautomers, racemics, prodrugs, metabolites thereof, or a pharmaceutically acceptable salt and/or solvate thereof, wherein X₁ and X₂ are -OMe, wherein R₁ and R₂ are -H, wherein X₃ or X'₃ are independently selected from the group ~~comprising~~consisting of hydrogen, glucosyl, fructosyl, galactosyl, mannosyl, fucosyl, isomaltosyl, maltosyl, cellobiosyl, gentiobiosyl, melibiosyl, palatinosyl, lactulosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, 2-amino-2-deoxy-glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy-galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-acetamido-2-deoxy-4-O-β-D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O-β-D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminyl lactosyl, 2-acetamido-2-deoxy-3-O-α-L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy-β-D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O-β-D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O-β-D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O-β-D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form ~~thereof~~thereof, pyranose or furanose form thereof, a disaccharide or a trisaccharide thereof, wherein X₄ is hydrogen, wherein X₅ and X₆ participates ~~to in~~a double bond between the carbon atoms in position 5 and 6, wherein X₆ is -H, wherein X₇ is selected from the group ~~comprising~~consisting of glucosyl, fructosyl, galactosyl, mannosyl, fucosyl, isomaltosyl, maltosyl, cellobiosyl, gentiobiosyl, melibiosyl, palatinosyl, lactulosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, 2-amino-2-deoxy-glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy-galactosyl, 2-

acetamido-2-deoxy-galactosyl, 2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminyllactosyl, 2-acetamido-2-deoxy-3-O- α -L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy- β -D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O- β -D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O- β -D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form thereof, pyranose or furanose form thereof, a disaccharide or a trisaccharide thereof, wherein at least one of X₃ and X'₃ is a glycosyl moiety selected from the group consisting of glucosyl, fructosyl, galactosyl, mannosyl, ribosyl, ribulosyl, xylulosyl, erythrosyl, erythrulosyl, rhamnosyl, threosyl, sorbosyl, psicose, tagatosyl, fucosyl, arabinosyl, xylofuranosyl, lyxosyl, talosyl, psicose, idosyl, gulosyl, altrosyl, allosyl, mannoheptulosyl, sedoheptulosyl, abequosyl, isomaltosyl, kojibiosyl, laminaribiosyl, nigerosyl, primeverosyl, rutinose, tyvelosyl, maltosyl, lactosyl, sucrosyl, cellobiosyl, trehalosyl, gentiobiosyl, melibiosyl, turanosyl, sophorosyl, isosucrosyl, raffinose, palatinosyl, lactulosyl, gentianosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, isomaltotriosyl, maltopentaosyl, maltohexaosyl, maltoheptaosyl, sicosyl, panosyl, isopanosyl, inosyl, N-acetylglucosaminyllactosyl, mannotriosyl, globotriosyl, erlosyl, neotrehalosyl, chitobiosyl, chitobiosemannosyl, glucosaminyllactosyl, N-acetyl-glucosaminyllactosyl, octylglucopyranosyl, octylribofuranosyl, cyclohexylglucopyranosyl, cyclohexylxylofuranosyl, benzylglucopyranosyl, benzylarabinofuranosyl, N-acetyl-lactosaminyllactosyl, acosaminyllactosyl, amictosyl, amylosyl, apiosyl, arcanosyl, ascarylosyl, bacillosaminyllactosyl, boivinosyl, cellotriosyl, chacotriosyl, chalcosyl, cladinosyl, colitosyl, cymarosyl, daunosaminyllactosyl, desosaminyllactosyl, D-glycero-L-gulo-heptosyl, diginosyl, digitalosyl, digitoxosyl, evalosyl, evernitrosyl, forosaminyllactosyl, fucosaminyllactosyl, garosaminyllactosyl, hamamelosyl, isolevoglucosenonyllactosyl, kanosaminyllactosyl, kansosaminyllactosyl, lactosaminyllactosyl, lactosediaminyllactosyl, fucitolyl, maltulosyl, mannosaminyllactosyl, melezitoyl, mycaminosyl, mycarosyl, mycinosyl, mycosaminyllactosyl, noviosyl, oleandrosyl, paratosyl, perosaminyllactosyl, planteosyl, pneumosaminyllactosyl, purpurosaminyllactosyl, quinovosaminyllactosyl, quinovosyl, rhamnitoyl, rhamnosaminyllactosyl, rhodinosyl, rhodosaminyllactosyl, sarmentosyl, solatriosyl, stachyosyl, streptosyl, umbelliferosyl, trehalosaminyllactosyl, 1,6-anhydro-D-glucopyranosyl, 1-hydroxy- α -

D-allopyranosyl, 2,3:5,6-di-O-isopropylidene-D-mannofuranosyl, 2-amino-2-deoxy-D-galactitolyl, 2-deoxyribosyl, 2-deoxyglucosyl, 5-amino-5-deoxy-D-glucopyranosyl, 6-deoxy-D-galactitolyl, 2-amino-2-deoxy glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-amino-2-deoxy mannosyl, 2-acetamido-2-deoxy-mannosyl, 2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminylactosyl, 2-acetamido-2-deoxy-3-O- α -L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy- β -D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O- β -D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O- β -D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form thereof, pyranose or furanose form thereof, combination thereof, deoxy derivatives thereof, hydroxyl-protected acetate or benzoyl derivatives thereof, amino derivatives thereof, amido derivatives thereof, thio derivatives thereof, di-, tri-, oligo- and polysaccharide thereof optionally substituted by one or more substituents independently selected from the group consisting of alkyl, aralkyl, aryl, Het¹, Het², cycloalkyl, alkyloxy, alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(alkyl)aminocarbonyl, aminosulfonyl, alkylS(=O)_t, hydroxy, cyano, halogen or amino optionally mono- or disubstituted wherein the substituents are independently selected from the group consisting of alkyl, aryl, aralkyl, aryloxy, arylamino, arylthio, aryloxyalkyl, arylaminoalkyl, aralkoxy, alkylthio, alkoxy, aryloxyalkoxy, arylaminoalkoxy, aralkylamino, aryloxyalkylamino, arylaminoalkylamino, arylthioalkoxy, arylthioalkylamino, aralkylthio, aryloxyalkylthio, arylaminoalkylthio, arylthioalkylthio, alkylamino, cycloalkyl and cycloalkylalkyl as indicated above and wherein n is 0.

11. (Currently amended) ~~A~~ The compound according to ~~any of claims 1 to 4~~ claim 1, stereoisomers, tautomers, racemics, prodrugs, metabolites thereof, or a pharmaceutically acceptable salt and/or solvate thereof, wherein X₁ and X₂ are -OMe, wherein R₁ and R₂ are -H, wherein X₃ participates together with X'₃ ~~to in an~~ in an oxo functional group or are independently selected from the group ~~comprising~~ consisting of hydrogen, hydroxyl, alkyloxy, wherein X₄ is selected from the group ~~comprising~~ consisting of glucosyl, fructosyl, galactosyl, mannosyl, fucosyl, isomaltosyl,

maltosyl, cellobiosyl, gentiobiosyl, melibiosyl, palatinosyl, lactulosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, 2-amino-2-deoxy-glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy-galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminylactosyl, 2-acetamido-2-deoxy-3-O- α -L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy- β -D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O- β -D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O- β -D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form thereof, pyranose or furanose form thereof, a disaccharide or a trisaccharide thereof, wherein X_5 and X_6 participates to in a double bond between the carbon atoms in position 4 and 5, wherein X_6 is -H, wherein X_7 is selected from the group comprisingconsisting of glucosyl, fructosyl, galactosyl, mannosyl, fucosyl, isomaltosyl, maltosyl, cellobiosyl, gentiobiosyl, melibiosyl, palatinosyl, lactulosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, 2-amino-2-deoxy glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy-galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminylactosyl, 2-acetamido-2-deoxy-3-O- α -L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy- β -D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O- β -D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O- β -D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form thereof, pyranose or furanose form thereof, a disaccharide or a trisaccharide thereof, and wherein n is 0.

12. (Currently amended) A ~~The~~ compound according to ~~any of claims 1 to 4~~ claim 1, stereoisomers, tautomers, racemics, prodrugs, metabolites thereof, or a pharmaceutically acceptable salt and/or solvate thereof, wherein X_1 and X_2 are -OMe, wherein R_1 and R_2 are -H, wherein X_3 or X'_3 are independently selected from the group comprisingconsisting of hydrogen, glucosyl, fructosyl, galactosyl, mannosyl, fucosyl,


isomaltosyl, maltosyl, cellobiosyl, gentiobiosyl, melibiosyl, palatinosyl, lactulosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, 2-amino-2-deoxy-glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy-galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-acetamido-2-deoxy-4-O-b-D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O-b-D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminylactosyl, 2-acetamido-2-deoxy-3-O-a-L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy-b-D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O-b-D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O-b-D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O-b-D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form thereof, pyranose or furanose form thereof, a disaccharide or a trisaccharide thereof, wherein X_4 is selected from the group ~~comprising~~consisting of glucosyl, fructosyl, galactosyl, mannosyl, fucosyl, isomaltosyl, maltosyl, cellobiosyl, gentiobiosyl, melibiosyl, palatinosyl, lactulosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, 2-amino-2-deoxy-glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy-galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-acetamido-2-deoxy-4-O-b-D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O-b-D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminylactosyl, 2-acetamido-2-deoxy-3-O-a-L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy-b-D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O-b-D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O-b-D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O-b-D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form thereof, pyranose or furanose form thereof, a disaccharide or a trisaccharide thereof, wherein X_5 and X_6 participates to ~~in~~a double bond between the carbon atoms in position 4 and 5, wherein X_6 is -H, wherein X_7 is selected from the group ~~comprising~~consisting of glucosyl, fructosyl, galactosyl, mannosyl, fucosyl, isomaltosyl, maltosyl, cellobiosyl, gentiobiosyl, melibiosyl, palatinosyl, lactulosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, 2-amino-2-deoxy glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy-galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-acetamido-2-deoxy-4-O-b-D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O-b-D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminylactosyl, 2-acetamido-2-deoxy-3-O-a-

L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy-b-D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O-b-D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O-b-D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O-b-D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form thereof, pyranose or furanose form thereof, a disaccharide or a trisaccharide thereof, wherein at least one of X_3 and X'_3 is a glycosyl moiety selected from the group consisting of glucosyl, fructosyl, galactosyl, mannosyl, ribosyl, ribulosyl, xylulosyl, erythrosyl, erythrulosyl, rhamnosyl, threosyl, sorbosyl, psicose, tagatosyl, fucosyl, arabinosyl, xylofuranosyl, lyxosyl, talosyl, psicose, idosyl, gulose, altriosyl, allosyl, mannoheptulosyl, sedoheptulosyl, abequosyl, isomaltosyl, kojibiosyl, laminaribiosyl, nigerosyl, primeverosyl, rutinosyl, tyvelosyl, maltosyl, lactosyl, sucrosyl, cellobiosyl, trehalosyl, gentiobiosyl, melibiosyl, turanosyl, sophorosyl, isosucrosyl, raffinose, palatinosyl, lactulosyl, gentianosyl, 3-mannobiosyl, 6-mannobiosyl, 3-galactobiosyl, 4-galactobiosyl, maltotriosyl, maltotetraosyl, isomaltotriosyl, maltopentaosyl, maltohexaosyl, maltoheptaosyl, sicosyl, panosyl, isopanosyl, inosyl, N-acetyl-galactosaminyl, mannotriosyl, globotriosyl, erlosyl, neotrehalosyl, chitobiosyl, chitobiosemannosyl, glucosaminyl, N-acetyl-glucosaminyl, octylglucopyranosyl, octylribofuranosyl, cyclohexylglucopyranosyl, cyclohexylxylofuranosyl, benzylglucopyranosyl, benzylarabinofuranosyl, N-acetyl-lactosaminyl, acosaminyl, amicetosyl, amylosyl, apiosyl, arcanosyl, ascarylosyl, bacillosaminyl, boivinosyl, cellotriosyl, chacotriosyl, chalcose, cladinosyl, colitosyl, cymarosyl, daunosaminyl, desosaminyl, D-glycero-L-gulo-heptosyl, diginosyl, digitalosyl, digitoxosyl, evalosyl, evernitrosyl, forosaminyl, fucosaminyl, garosaminyl, hamamelosyl, isolevoglucosenonyl, kanosaminyl, kansosaminyl, lactosaminyl, lactosediaminyl, fucitolyl, maltulosyl, mannosaminyl, melezitose, mycaminosyl, mycarosyl, mycinosyl, mycosaminyl, noviosyl, oleandrosyl, paratosyl, perosaminyl, planteosyl, pneumosaminyl, purpurosaminyl, quinovosaminyl, quinovosyl, rhamnitolyl, rhamnosaminyl, rhodinosyl, rhodosaminyl, sarmentosyl, solatriosyl, stachyosyl, streptosyl, umbelliferosyl, trehalosaminyl, 1,6-anhydro-D-glucopyranosyl, 1-hydroxy- α -D-allopyranosyl, 2,3:5,6-di-O-isopropylidene-D-mannofuranosyl, 2-amino-2-deoxy-D-galactitolyl, 2-deoxyribosyl, 2-deoxyglucosyl, 5-amino-5-deoxy-D-glucopyranosyl, 6-deoxy-D-galactitolyl, 2-amino-2-

deoxy glucosyl, 2-acetamido-2-deoxy-glucosyl, 2-amino-2-deoxy galactosyl, 2-acetamido-2-deoxy-galactosyl, 2-amino-2-deoxy mannosyl, 2-acetamido-2-deoxy-mannosyl, 2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 2-amino-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, 6'-N-acetylglucosaminyllactosyl, 2-acetamido-2-deoxy-3-O- α -L-fucosyl-D-glucosyl, 6-O(2-acetamido-2-deoxy- β -D-glucosyl)-D-galactosyl, 2-acetamido-2-deoxy-3-O- β -D-galactosyl-D-glucosyl, 2'-acetamido-2'-deoxy-3-O- β -D-glucosyl-D-galactosyl, 3-fucosyl-D-lactosyl, 3-fucosyl-2-acetamido-2-deoxy-4-O- β -D-galactosyl-D-glucosyl, L or D isomers thereof, α or β form thereof, pyranuronic or furanuronic form thereof, pyranose or furanose form thereof, combination thereof, deoxy derivatives thereof, hydroxyl-protected acetate or benzoyl derivatives thereof, amino derivatives thereof, amido derivatives thereof, thio derivatives thereof, di-, tri-, oligo- and polysaccharide thereof optionally substituted by one or more substituents independently selected from the group consisting of alkyl, aralkyl, aryl, Het¹, Het², cycloalkyl, alkyloxy, alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(alkyl)aminocarbonyl, aminosulfonyl, alkylS(=O)_t, hydroxy, cyano, halogen or amino optionally mono- or disubstituted wherein the substituents are independently selected from the group consisting of alkyl, aryl, aralkyl, aryloxy, arylamino, arylthio, aryloxyalkyl, arylaminoalkyl, aralkoxy, alkylthio, alkoxy, aryloxyalkoxy, arylaminoalkoxy, aralkylamino, aryloxyalkylamino, arylaminoalkylamino, arylthioalkoxy, arylthioalkylamino, aralkylthio, aryloxyalkylthio, arylaminoalkylthio, arylthioalkylthio, alkylamino, cycloalkyl and cycloalkylalkylas indicated above and wherein n is 0.

13. (Currently amended) ~~Compound~~ A compound of formula I, stereoisomers, tautomers, racemics, prodrugs, metabolites thereof, or a pharmaceutically acceptable salt and/or solvate thereof, wherein X₁, X₂, X₃, X'₃, X₄, X₅, X₆, X₇, R₁, R₂ and n are selected as indicated in Table A.


14. (Currently amended) ~~Method~~ A method for synthesizing a compound ~~having comprising~~ the structural formula I



formula I

wherein X_1 , X_2 , X_3 , X_4 , X_5 , X_6 , X_7 , R_1 , R_2 and n are independently selected from the group as indicated in ~~any of claims 1 to 13~~ claim 1, said method comprising the steps of

a) providing a starting material ~~having~~ comprising the structural formula IV,




formula IV

wherein X_3 participates together with X'_3 ~~to in~~ in an oxo functional group, or wherein X_3 and X'_3 are independently selected from the group ~~comprising~~ consisting of hydrogen, hydroxyl, sulfur, oxyalkyl, oxycarbonyl, alkyl, Het¹alkyl, alkyloxycarbonyl, alkenyl, alkynyl, aminoalkyl, aminoacyl, alkylcarbonylamino, alkylthiocarbonylamino, Het¹, optionally substituted by one or more substituents independently selected from the group ~~comprising~~ consisting of alkyl, aralkyl, aryl, Het¹, Het², cycloalkyl, alkyloxy, alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(alkyl)aminocarbonyl, aminosulfonyl, alkylS(=O)t, hydroxy, cyano, halogen or amino optionally mono- or disubstituted wherein the substituents are independently selected from the group ~~comprising~~ consisting of alkyl, aryl, aralkyl, aryloxy, arylamino, arylthio, aryloxyalkyl, arylaminoalkyl, aralkoxy, alkylthio, alkoxy, aryloxyalkoxy, ~~arylaminealkoxy~~ arylaminealkoxy, aralkylamino, aryloxyalkylamino, arylaminoalkylamino, arylthioalkoxy, arylthioalkylamino, aralkylthio, aryloxyalkylthio, arylaminoalkylthio, arylthioalkylthio, alkylamino, cycloalkyl and cycloalkylalkyl;

wherein X_7 is selected from the group ~~comprising~~consisting of hydrogen, oxygen, halogen, oxo, carbonyl, thiocarbonyl, hydroxyl, alkyl, aryl, Het¹, Het¹alkyl, Het¹aryl, alkenyl, alkynyl, hydroxyalkyl, hydroxycarbonyl, hydroxycarbonylalkyl, hydroxycarbonylaryl, hydroxycarbonyloxyalkyl optionally substituted by one or more substituents independently selected from the group ~~comprising~~consisting of alkyl, aralkyl, aryl, Het¹, Het², cycloalkyl, alkyloxy, alkyloxycarbonyl, carboxyl, aminocarbonyl, mono- or di(alkyl)aminocarbonyl, aminosulfonyl, alkylS(=O)t, hydroxy, cyano, halogen or amino optionally mono- or disubstituted wherein the substituents are independently selected from the group ~~comprising~~consisting of alkyl, aryl, aralkyl, aryloxy, arylamino, arylthio, aryloxyalkyl, arylaminoalkyl, aralkoxy, alkylthio, alkoxy, aryloxyalkoxy, arylaminealkoxyarylaminealkoxy, aralkylamino, aryloxyalkylamino, arylaminoalkylamino, arylthioalkoxy, arylthioalkylamino, aralkylthio, aryloxyalkylthio, arylaminoalkylthio, arylthioalkylthio, alkylamino, cycloalkyl and cycloalkylalkyl; and wherein P is a protecting group,


b) effecting reaction between the compound of step a) with an organometallic compound ~~having~~comprising the structural formula V



formula V

wherein X_1 , X_2 , R_1 , R_2 and n are independently selected from the group as indicated in ~~any of claims 1 to 13~~claim 1, wherein W is a metal or a combination of metals and wherein Hal is a halogen atom,

to result in an intermediate ~~having~~comprising the structural formula III'



formula III'

wherein X_1 , X_2 , R_1 , R_2 and n are independently selected from the group as indicated in any of claims 1 to 13, wherein X_3 , X'_3 , X_7 are independently selected from the group as indicated in step a) and wherein P is a protecting group,


c) effecting reaction between the compound of step b) with an organometallic compound having comprising the structural formula VI



formula VI

wherein X'_3 is selected from the group as indicated in step a), wherein W is a metal or a combination of metals, and wherein Hal is a halogen atom,


to result in an intermediate having comprising the structural formula III



formula III

wherein X_1 , X_2 , R_1 , R_2 and n are independently selected from the group as indicated in ~~any of claims 1 to 13~~ claim 1, wherein X_3 , X'_3 , X_7 are independently selected from the group as indicated in step a), wherein P is a protecting group,

d) deprotecting the X_7 group of the compound obtained in step c) to form an compound having comprising the structural formula II



formula II

wherein X_1 , X_2 , R_1 , R_2 and n are independently selected from the group as indicated in ~~any of claims 1 to 13~~ claim 1, wherein X_3 , X'_3 , X_7 are independently selected from the group as indicated in step a), and

e) coupling an O-protected glycosyl or non-protected glycosyl to form a compound of formula IIB wherein X_1 , X_2 , R_1 , R_2 and n are independently selected from the group as indicated in ~~any of claims 1 to 13~~ claim 1, wherein X_3 and X'_3 are independently selected from the group as indicated in step a), and X_7 is an O-protected glycosyl or a non-protected glycosyl, and

f) deprotecting the O-protected groups of glycosyl to form a compound of formula IB wherein X_1 , X_2 , X_4 , X_5 , X_6 , R_1 , R_2 and n are independently selected from the group as indicated in ~~any of claims 1 to 13~~ claim 1, wherein X_3 , X'_3 are independently selected from the group as indicated in step a), and X_7 is selected from the group ~~comprising~~ consisting of glycosyl, thio derivatives thereof, amino derivatives thereof, amido derivatives thereof, hydroxyl-protected derivatives thereof.


15. (Currently amended) ~~Method~~ The method according to claim 14, wherein step e) consists of reacting the compound of step d) with an oxidizing reagent to form an intermediate and reducing said intermediate with a reducing reagent to result in another intermediate ~~having~~ comprising the structural formula I wherein X_1 , X_2 , R_1 , R_2 and n are independently selected from the group as indicated in ~~any of claims 1 to 13~~, and X_3 or X'_3 , and X_4 and X_7 are hydroxyl and continuing the reaction with steps e) and f) according to claim 14 to form a glycosylated steroid compound of structural formula I.

16. (Currently amended) ~~Method~~ The method according to claim 14, wherein step c) consists of reacting the compound of step b) with an O-protected glycosyl or non-

protected glycosyl to result in an intermediate having comprising the structural formula III wherein ~~X₁, X₂, R₁, R₂ and n are independently selected from the group as indicated in any of claims 1 to 13, wherein X₃, X₇ are independently selected from the group as indicated in step a) of claim 14, wherein P is a protecting group, and wherein X₃ or X'₃ is an O-protected glycosyl or a non-protected glycosyl and continuing the reaction with steps d), e) and f) according to claim 14 to form a glycosylated steroid compound of structural formula I.~~


17. (Currently amended) A compound obtainable by any of the steps according to the method of ~~any of claims 14 to 16~~ claim 14.

18. (Currently amended) ~~A~~ The compound according to claim 1, of structural formula:




herein designated as compound **UBS3268**

19. (Currently amended) ~~A~~ The compound according to claim 1, of structural formula:




herein designated as compound **UBS3270**

20. (Currently amended) ~~A~~ The compound according to claim 1, of structural formula:




herein designated as compound **UBS3285**

21. (Currently amended) A-The compound according to claim 1, of structural formula:




herein designated as compound **UBS3327**

22. (Currently amended) A-The compound according to claim 1, of structural formula:




herein designated as compound **UBS3328**

23. (Currently amended) A-The compound according to claim 1, of structural formula:




herein designated as compound **UBS3501**

24. (Currently amended) A-The compound according to claim 1, of structural formula:




herein designated as compound **UBS3585**

25. (Currently amended) A The compound according to claim 1, of structural formula:




herein designated as compound **UBS3597**

26. (Currently amended) A The compound according to claim 1, of structural formula:




herein designated as compound **UBS3976**

27. (Currently amended) A The compound according to claim 1, of structural formula:




herein designated as compound **UBS4066**

28. (Currently amended) A The compound according to claim 1, of structural formula:




herein designated as compound **UBS4067**

29. (Currently amended) A-The compound according to claim 1, of structural formula:




herein designated as compound **UBS4095**

30. (Currently amended) A-The compound according to claim 1, of structural formula:




herein designated as compound **UBS4104**

31. (Currently amended) A-The compound according to claim 1, of structural formula:




herein designated as compound **UBS4109**

32. (Currently amended) A-The compound according to claim 1, of structural formula:



herein designated as compound **UBS4209**

33. (Currently amended) A The compound according to claim 1, of structural formula:



herein designated as compound **UBS4373**

34. (Cancelled)

35. (Cancelled)

36. (Currently amended) ~~Use of~~ A method for treating cancer which comprises administering a compound according to any of claims 1 to 13 and 17 to 33 claim 1 to an individual in need thereof for the preparation of a medicament for treating cancer.

37. (Currently amended) A pharmaceutical composition comprising a pharmaceutically acceptable excipient and a therapeutically effective amount of a compound according to ~~any of claims 1 to 13 and 17 to 33~~ claim 1.

38. (Cancelled)

39. (Currently amended) ~~Method~~ A method of treating cancer comprising administrating to an individual in need of such treatment a pharmaceutical composition according to claim 37.